# **Moth Trapping Report**

## **Summary**

Location:	Beachampton (SP7736)	
Date:	15/05/2024	
Trap(s):	MV Robinson Trap	
Recorders(s):	Chris Chapman	
Moth Count:	47	
Species Count: (see notes)	19 New for Year: 19 New for Location: 5 Total new species for Location this Year: 5 Species Count for Year: 19 Species Total for Location: 81	
Non-species Count:	0	

## **Full List**

#### NFG = New Species For Location/Garden

Species	Count	Type	Comment
Brimstone Moth (Opisthograptis luteolata)	1	Macro	A common, resident species. Double brooded, flying from late April to mid July and then again from early August to early October. The larvae feed on hawthorn, apple, plum, hazel and sloe.
Common Marble (Celypha lacunana)	1	Micro	Resident. Larvae feed in the young shoots of almost any shrubs and plants, including Epilobium and Hypericum species.
Figure of Eighty (Tethea ocularis)	2	Macro	
Flame Shoulder (Ochropleura plecta)	2	Macro	Resident. Larvae feed on various unidentified wild and cultivated plants that have been listed from elsewhere.
Green Carpet (Colostygia pectinataria)	7	Macro	Resident. Larvae feed on bedstraw.
Herald (Scoliopteryx libatrix) NFG	1	Macro	Resident. Larvae feed on sallow and other willows.
Muslin Moth (Diaphora mendica) NFG	2	Macro	A resident species. Males are widespread and usually occur sparingly in a wide range of habitats. Females are comparatively scarce and largely confined to woodland clearings, and are only seen flying by day. Single-brooded, flying mainly from early April
Pale Tussock (Calliteara pudibunda)	1	Macro	
Plum Tortrix (Hedya pruniana)	1	Micro	Resident. Larvae feed on the young shoots of sloe and plum.
Poplar Hawk-moth (Laothoe populi)	3	Macro	Resident . Larvae feed on goat willow, grey willow, aspen, poplars, sallows and occasionally beech.
Red Twin-spot Carpet (Xanthorhoe spadicearia) NFG	1	Macro	
Rustic Shoulder-knot (Apamea sordens)	8	Macro	
Sandy Carpet (Perizoma flavofasciata) NFG	1	Macro	
Scorched Wing (Plagodis dolabraria)	1	Macro	
Swallow Prominent (Pheosia tremula) NFG	1	Macro	A resident occurring singly or sparingly in a wide range of habitats.

Species	Count	Туре	Comment
			Probably generally distributed. Double-brooded, flying mainly from late April to mid June and again from late July to early September. The larvae feed on sallow, poplar and aspen.
Treble Lines (Charanyca trigrammica)	10	Macro	
Turnip Moth (Agrotis segetum)	1	Macro	
White Ermine (Spilosoma lubricipeda)	2	Macro	A resident and occasional immigrant. Previously single-brooded, flying from late April to mid July but, incresasingly, from mid August to late September. Larvae feed on ivy flowers, dock, plantain, bindweed, hop, goosefoot and bedstraw
Yellow-faced Bell (Notocelia cynosbatella)	1	Micro	Resident. The larvae feed in the shoots of wild and cultivated roses.

## **Images**



Green Carpet (Colostygia pectinataria)



Herald (Scoliopteryx libatrix)



Muslin Moth (Diaphora mendica)



Sandy Carpet (Perizoma flavofasciata)



Swallow Prominent (Pheosia tremula)



Treble Lines (Charanyca trigrammica)

### **Notes**

The numbers in the **Species Count** section of the **Summary** section only include moths that have been identified to species level.

You can see more details of each species from the UK Moths website by clicking on the name of the species. This doesn't apply to records at genus level or aggregates.

Produced by Chris Chapman (Plumpton & East Chiltington Wildlife Group)