



Moth Trapping Report

Summary

Location:	Stoke-sub-Hamdon (ST4817)
Date:	24/10/2025
Trap(s):	MV Robinson Trap
Recorders(s):	Chris Chapman
Moth Count:	64
Species Count:	16
Non-species Count:	1

Full List

Species	Count	Type	Comment
Barred Sallow (<i>Tiliacea aurago</i>)	3	Macro	Resident. The larval foodplants are maple and beech.
Beaded Chestnut (<i>Agrochola lychnidis</i>)	20	Macro	Resident. The larvae feed on unspecified grasses and low growing plants.
Black Rustic (<i>Aporophyla nigra</i>)	5	Macro	Resident. Larvae feed on heather, petty whin, dock, dandelion and foxglove.
Brown-spot Flat-body (<i>Agonopterix alstromeriana</i>)	1	Micro	
Common Marbled Carpet (<i>Dysstroma truncata</i>)	1	Macro	Resident. Larvae have a wide variety of food plants.
Feathered Thorn (<i>Colotois pennaria</i>)	1	Macro	
Green-brindled Crescent (<i>Allophyes oxyacanthae</i>)	9	Macro	Resident. Larvae feed on hawthorn, sloe and garden apple.
Large Yellow Underwing (<i>Noctua pronuba</i>)	2	Macro	Resident reinforced by regular immigration. Larvae feed on unspecified garden vegetables and flowers.
Merveille du Jour (<i>Griposia aprilina</i>)	4	Macro	Resident. Larvae feed on oak.
November Moth agg. (<i>Epirrita dilutata</i> agg.)	1	Micro	
Red-line Quaker (<i>Leptologia lota</i>)	3	Macro	Resident. The larvae feed on Sallow and willow (including grey willow).
Rush Veneer (<i>Nomophila noctuella</i>)	3	Micro	Immigrant appearing each year in widely varying numbers. Larvae feed on clover and unspecified grasses.
Sallow (<i>Cirrhia icteritia</i>)	1	Macro	
Setaceous Hebrew Character (<i>Xestia c-nigrum</i>)	1	Macro	Resident species reinforced by episodic immigration. Larvae feed on dock and probably other herbaceous plants.
Vine's Rustic (<i>Hoplodrina ambigua</i>)	2	Macro	Resident and immigrant species. The larval foodplant has not been recorded in Sussex.
White-point (<i>Mythimna albipuncta</i>)	6	Macro	Immigrant and episodic resident. Larvae feed on unspecified grasses.
Yellow-line Quaker (<i>Leptologia macilentata</i>)	1	Macro	Resident. The larvae feed on oak, beech and hawthorn.

Images



Barred Sallow (*Tiliacea aurago*)



Beaded Chestnut (*Agrochola lychnidis*)



Black Rustic (*Aporophyla nigra*)



Feathered Thorn (*Colotois pennaria*)



Merveille du Jour (*Griposia aprilina*)



Red-line Quaker (*Leptologia lota*)



Setaceous Hebrew Character (*Xestia c-nigrum*)



Yellow-line Quaker (*Leptologia macilenta*)

Notes

You can see more details of each species from the [UK Moths website](#) by clicking on the name of the species. This doesn't apply to records at genus level or aggregates.

Produced by Chris Chapman ([Plumpton & East Chiltington Wildlife Group](#))