



Moth Survey Report

Summary

Location:	Kastel House, Veranda ()
Date:	03/06/2026
Survey Method(s):	Attracted to a lighted Window
Recorders(s):	Chris Chapman
Moth Count:	29
Species Count: (see notes)	17 New for Year: 10 New for Location: 5 Total new species for Location this Year: 13 Species Count for Year: 23 Species Total for Location: 216
Non-species Count:	1

Full List

NFG = New Species For Location/Garden NFY = New Species For Year

Species	Count	Type	Comment
Delicate (<i>Mythimna vitellina</i>) NFY	1	Macro	Immigrant and suspected episodic resident species. Larvae feed on various grasses.
Druid (<i>Aedia funesta</i>) NFY NFG	1	Macro	
Four-spotted (<i>Tyta luctuosa</i>) NFY NFG	2	Macro	
Grass Wave (<i>Perconia strigillaria</i>)	1	Macro	
Horisme corticata (<i>Horisme corticata</i>)	2	Macro	
<i>Idaea deversaria</i> (<i>Idaea deversaria</i>) * NFY	1	Macro	
Lace Border (<i>Scopula ornata</i>) NFY	1	Macro	
Marbled Minor agg. (<i>Oligia strigilis</i> agg.)	4	Macro	
Meadow Grey (<i>Scoparia pyralella</i>)	2	Micro	
Portland Ribbon Wave (<i>Idaea degeneraria</i>)	4	Macro	An increasingly common immigrant.
Rusty-dot Pearl (<i>Udea ferrugalis</i>) NFY	1	Micro	Immigrant. Larvae feed on hemp agrimony, strawberry and woundwort.
Spotted Sulphur (<i>Acontia trabealis</i>)	1	Macro	
Tawny Wave (<i>Scopula rubiginata</i>) NFY	1	Macro	
Triangle-marked Roller (<i>Ancylis achatana</i>) NFY NFG	1	Micro	
Vine's Rustic (<i>Hoplodrina ambigua</i>) NFY	1	Macro	Resident and immigrant species. The larval foodplant has not been recorded in Sussex.
White Ermine (<i>Spilosoma lubricipeda</i>) NFY NFG	1	Macro	A resident and occasional immigrant. Previously single-brooded, flying from late April to mid July but, increasingly, from mid August to late September. Larvae feed on ivy flowers, dock, plantain, bindweed, hop, goosefoot and bedstraw
White-point (<i>Mythimna albipuncta</i>)	3	Macro	Immigrant and episodic resident. Larvae feed on unspecified

Species	Count	Type	Comment
			grasses.
Willow Beauty (Peribatodes rhomboidaria) NFY	1	Macro	Redsident. Larvae feed on privet, lilac, ivy, honeysuckle, hawthorn, dock, heather and gorse.

Images



Druid (Aedia funesta)



Four-spotted (Tyta luctuosa)



Horisme corticata (Horisme corticata)



Lace Border (*Scopula ornata*)



Spotted Sulphur (*Acontia trabealis*)



Tawny Wave (*Scopula rubiginata*)



Triangle-marked Roller (*Ancyliis achatana*)

Notes

The numbers in the **Species Count** section of the **Summary** section only include moths that have been identified to species level.

An * by the name of the species indicates that the species has not been verified by the County Recorders yet. In total this applies to 1 records for this survey.

You can see more details of each species from the [UK Moths website](#) by clicking on the name of the species. This doesn't apply to records at genus level or aggregates.

For further details of the records see [Green Cross Nature](#)

